

The true solution for rapid diagnostics of acute infections















mariPOC® optimizes hospital processes, improves patient care and reduces unnecessary antibiotic use!



1. Examine the patient



2. Select the applicable mariPOC® test



3. Take a sample



 Run an automatic mariPOC® analysis for use in evidence-based medicine mariPOC® is an automated test system for the rapid multianalyte identification of acute infectious disease pathogens. The test system can be used in all areas of patient care. mariPOC® combines specific gene expression testing with unique laser-based TPX detection technology. The innovative mariPOC® technology's advantages include superior sensitivity, specificity and efficiency.

The technology includes sophisticated lab-level autoverification of sample analysis to ensure utmost reliability. The intuitive user interface displays the results of the automated fluorescence reading, and the system enables bidirectional connection to a LIS. The test system provides qualitative or quantitative results depending on the user needs.

mariPOC® is a walk-away device that is very easy to use. The device can be operated after a short introduction. The short hands-on time is 1 to 3 minutes depending on the test being performed.


Why choose mariPOC®?

Cost-efficient solution

Rapid results minimize delays in diagnosis and reduce the turnaround time.

Sensitive & Specific

mariPOC® provides results with central laboratory accuracy near the patient.



High throughput

mariPOC® enables the continuous feed of new samples, allowing the analysis of more than 50 samples during a single work shift!

Flexible test plate change

It is possible to run multiple applications on a single device.

LIS connectability

mariPOC® can be connected to a LIS for ease of use and to prevent human errors.

Automated analysis and result reporting

This feature ensures truly reliable results with every measurement.

Performance

Test	mariPOC® compared to	Sensitivity	Specificity
	PCR		
	SARS coronavirus 2	92.3%	100%
Respi+	Influenza A virus	92.3%	99.8%
кезріт	Influenza B virus	87.5%	100%
	RSV	88.6%	100%
	Bacterial culture		
Pharyn	Group A streptococci	Pharyn 150% Quick StrepA 100%	100%
	Immunochromatography		
CDI	C. difficile GDH	100%	98.8%
	C. difficile toxins A/B	> 100%	100%



Thomas E. et al. 29th ECCMID 2019, Amsterdam, Netherlands. Abstract and poster #P0114.

Sanbonmatsu-Gámez S. et al. (2015) Diagn Microbiol Infect Dis. 83:252-256. Ivaska L. et al. (2013) J Clin Virol. 57:136-140.

Vakkila J. et al. (2015) J Clin Microbiol. 53:2079-2083.



Respi+

SARS coronavirus 2
Influenza A virus
Influenza B virus
Respiratory syncytial virus
Human metapneumovirus
Human coronavirus OC43
Parainfluenza viruses 1, 2, 3
Adenovirus
Streptococcus pneumoniae

Quick Flu+

SARS coronavirus 2 Influenza A virus Influenza B virus Respiratory syncytial virus

Quick Flu/RSV

Influenza A virus Influenza B virus Respiratory syncytial virus



for pharyngitis

Quick StrepA

Group A streptococci

Pharyn

Group A streptococci Adenovirus



for gastroenteritis

Gastro

Norovirus GII.4 Norovirus GI Rotavirus Adenovirus Campylobacter spp.



for Clostridium difficile infection

CDI

C. *difficile* GDH C. *difficile* Toxins A & B

Ensure correct pathogen-specific diagnosis and treatment with mariPOC®'s broad test panels!

REF	Test plate	mariPOC® test	Tests/ plate
1184M	mariPOC® Respi+	Respi+	22
1194M	mariPOC® RTI combi+	Respi+, Quick Flu+ Pharyn, Quick StrepA	22 22
1204S	mariPOC® SARS-CoV-2	SARS-CoV-2	308
1124M	mariPOC® Pharyn	Pharyn, Quick StrepA	66
1214M	mariPOC® Quick combi+	Quick Flu+ Quick StrepA	22 22
2017M	mariPOC® Gastro	Gastro	44
2027M	mariPOC® Gastro CDI combi	Gastro CDI (GDH + Toxins A/B)	44 44
2037M	mariPOC® CDI	CDI (GDH + Toxins A/B)	44